

# Clinical Considerations for Managing Chronic Enteropathy Patients: Why Improving GI Microbiome Health Matters



**Linda Toresson, DVM, PhD**  
Evidensia Specialist Animal Hospital Helsingborg, Sweden and Department of Equine and Small Animal Internal Medicine, Helsinki University Finland

## Q The intestinal microbiome is a complex ecosystem. What comprises a healthy microbiome versus an unhealthy one?

**A** I like to describe the composition of the healthy intestinal microbiome as a “good party” and dysbiosis as a “bad party” that takes place in the gut. Consider the following characteristics of each:

Healthy microbiome (the “good party”)	Dysbiosis (the “bad party”)
Diverse “guests” (bacterial taxa) that buffer harmful bacteria	Less buffering against unwelcome “guests” (harmful bacteria)
Plenty of food for colonocytes in the form of short-chain fatty acids (SCFAs) produced by beneficial bacteria	No beneficial microbes producing SCFAs to feed colonocytes
Good, anti-inflammatory atmosphere rich in SCFAs, indole and secondary bile acids	Toxic, pro-inflammatory atmosphere due to the lack of beneficial microbes
No leakage thanks to protective SCFAs and indole	Intestinal leakage into the bloodstream
Healthy motility	Accelerated motility due to increased numbers of primary bile acids
Controlled proliferation of less-beneficial microbes	Proliferation of <i>E. coli</i> , <i>Clostridium perfringens</i> and <i>Clostridium difficile</i>

## Q Culture and sensitivity testing of faecal samples is commonly conducted with chronic enteropathy (CE) patients to determine which antibiotic will eliminate the suspected pathogen. What is problematic about this approach?

**A** Veterinarians gain limited information from conventional faecal cultures. Because strict anaerobes cannot be detected using standard cultivation methods, only around 5% of the bacterial species in the faecal microbiome can be identified. A recent study compared faecal culture results from healthy dogs versus from dogs with chronic diarrhoea at three different commercial laboratories. The cultures failed to distinguish between diseased and healthy dogs and two of the three labs reported an abnormal faecal microbiome in more healthy dogs than dogs with chronic diarrhoea.<sup>1</sup>

## Q Given the role of dysbiosis in dogs with CE, which tools can positively affect the intestinal microbiome?

**A** Studies have been conducted to evaluate various tools and their value in managing canine patients with CE.

- **Diet:** In a study of dogs with food-responsive enteropathy (FRE),

feeding a diet without animal protein resulted in a partial recovery of the faecal microbiota.<sup>2</sup>

- **Anti-inflammatory medication:** A study conducted on dogs with steroid-responsive enteropathy showed that steroid therapy eliminated clinical signs of disease after 8 weeks; however, normalisation of the faecal microbiome and metabolome was not noted until one year after treatment.<sup>3</sup>
- **Probiotics:** There has been some evidence that probiotics can reduce inflammation and foster the growth of beneficial bacteria such as *faecalibacterium*;<sup>4</sup> however, such results are probiotic-dependent and should not be generalised.
- **Prebiotics:** While most studies have been conducted in healthy animals, there is evidence that prebiotic fibres can improve stool quality and microbiome diversity.<sup>5</sup>
- **Faecal microbiota transplantation (FMT):** In one study, dogs with inflammatory bowel disease had a lower proportion of *Fusobacterium* (a beneficial bacterium that produces butyric acid, an SCFA) versus healthy dogs; however, this was altered with a single FMT.<sup>6</sup>
- **Antibiotics:** Antibiotics are not a good way to positively affect the microbiome. Broad-spectrum antibiotics such as metronidazole can create dysbiosis even in healthy dogs.<sup>7</sup> While dogs with CE may initially respond to both antibiotic and immunosuppressant (e.g., steroid) therapy,<sup>8</sup> a high percentage are likely to relapse within weeks when given antibiotics.<sup>9</sup>

## Q Are there other reasons that use of broad-spectrum antibiotics should be reconsidered in CE patients?

**A** Decisions to use antibiotics in companion animals are not without consequence. A recent study in Portugal looked at 20 households with humans, dogs and cats—all healthy. In 50% of the households, human/animal sharing of at least one gene for antimicrobial resistance (AMR) was documented.<sup>10</sup> A common belief is that tylosin is not used in humans and thus is safe for use in dogs; however, its administration is associated with AMR to erythromycin—a drug that is important to people.<sup>11</sup>

Antibiotic stewardship is an extremely important issue; as veterinarians, we must be careful about making decisions to use antibiotics in a patient. When we do, we should be aware that the decision could affect the pet’s family and advise them accordingly.



1. Werner M, Suchodolski JS, Lidbury JA, et al. Diagnostic value of fecal cultures in dogs with chronic diarrhea. *J Vet Int Med.* Vol 35, No 1, Jan/Feb 2021. 199-208.  
 2. Bresciani F, Minamoto Y, Suchodolski JS, et al. Effect of an extruded animal protein-free diet on fecal microbiota of dogs with food-responsive enteropathy. *J Vet Int Med.* 2018;32:1903-1910.  
 3. Pilla R, Guard BC, Blake AB, et al. Long-Term Recovery of the Fecal Microbiome and Metabolome of Dogs with Steroid-Responsive Enteropathy. *Animals.* 2021; 11(9):2498. <https://doi.org/10.3390/ani11092498>.  
 4. Rossi G, Pengo G, Caldin M, et al. Comparison of microbiological, histological and immunomodulatory parameters in response to treatment with either combination therapy with prednisone and metronidazole or probiotic VSL#3 strains in dogs with idiopathic inflammatory bowel disease. *PLoS One.* 2014 Apr 10;9(4):e94699.  
 5. Jackson MI, Jewell DE. Balance of saccharolysis and proteolysis underpins improvements in stool quality induced by adding a fibre bundle containing bound polyphenols to either hydrolyzed meat or grain-rich foods. *Gut Microbes.* 2019, Vol. 10, No. 3 298-320.  
 6. Niina A, Kibe R, Suzuki R, et al. Fecal microbiota transplantation as a new treatment for canine inflammatory bowel disease. *Biosci Microbiota Food Health.* 2021;40(2):98-104.  
 7. Pilla R, Gaschen FP, Barr JW, et al. Effects of metronidazole on the fecal microbiome and metabolome in healthy dogs. *J Vet Int Med.* 2020 Sep;34(5):1853-1866.  
 8. Treese JS, Nolte I, Bach JP. Chronic enteropathy in dogs—retrospective case analysis and results of an owner survey. *Tierärztl Prax Ausg K Kleintiere Heimtiere.* 2021 Apr;49(2):94-103.  
 9. Westermarck E, Skrzypczak T, Harmoinen J, et al. Tylosin-responsive chronic diarrhea in dogs. *J Vet Int Med.* Mar-Apr 2005;19(2):177-86.  
 10. Belas A, Menezes J, Gama LT, et al. Sharing of Clinically Important Antimicrobial Resistance Genes by Companion Animals and Their Human Household Members. *Microb Drug Resist.* 2020 Oct;26(10):1174-1185.  
 11. Jackson CR, Fedorka-Cray, PJ, Barrett JB, et al. Effects of Tylosin Use on Erythromycin Resistance in Swine. *Applied and Environmental Microbiology.* July 2004;4205-4210.


# Exploring the Gut Microbiome and its Role in Your Patients' Health

The gastrointestinal tract is primarily known as the system responsible for digestion and nutrient absorption. But it also plays a remarkable role in overall health—both for humans and pets.

## The gut is home to:

70%

of the immune cells, protecting the body from harmful (pathogens) or foreign substances (antigens).<sup>1</sup>



A complex microbial ecosystem with trillions of microorganisms, known as the gut microbiome

## How is the Microbiome Linked to Health?

The microbiome can impact a pet's overall health in many ways:<sup>2</sup>




## Why is Microbiome Balance So Important?

Because the microbiota in the gut help support overall health, it is important to maintain a balance between beneficial and potentially pathogenic bacteria. An imbalanced or maladapted microbiota is called dysbiosis.

### What can lead to dysbiosis?

Antibiotics	Illness
Stress	Diet change
Age	Weaning



### Dysbiosis is associated with a variety of health conditions:<sup>2</sup>

- IBD
- Obesity
- Cardiovascular disease
- Immune-mediated conditions
- Neurodevelopmental conditions

## How Can You Help? Promote Microbiome Balance with Nutrition



### PROBIOTICS

Live microorganisms that, when consumed in adequate amounts, confer a health benefit on the host.<sup>3</sup>

Probiotics are strain-specific and dose-dependent, meaning that different strains have different effects and they must be given in specific amounts.



### PREBIOTICS

Soluble fibers that are selectively fermented by beneficial bacteria, resulting in increased microbial diversity and the production of short-chain fatty acids that nourish colonocytes.<sup>4</sup>

Products that contain both prebiotics and probiotics are called synbiotics.



### COLOSTRUM

Contains bioactives and antibodies that initiate a beneficial immune response and help to stabilize intestinal microflora.<sup>5</sup>

Studies show that when fed diets supplemented with bovine colostrum bio-actives, both kittens<sup>6</sup> and adult dogs<sup>5</sup> have stronger responses to vaccinations.



SCAN ME

## Explore the gut microbiome... and beyond

What comprises the gut microbiome? How can the brain influence gut microbiota? What other microbiomes exist beyond the gut? There are many aspects to this topic and the Purina Institute has them covered. Just click on one of 10 areas to take a deeper dive into each. Visit <https://www.purinainstitute.com/microbiome-forum/microbiome-fundamentals> or scan the QR code to start exploring.

1. Vighi G, Marcucci F, Sensi L, et al. Allergy and the gastrointestinal system. *Clin Exp Immunol*. 2008 Sep;153 Suppl 1(Suppl 1):3-6.

2. Barco PC, McMichael MA, Swanson KA, et al. The Gastrointestinal Microbiome: A Review. *J Vet Intern Med*. 2018. Jan;32(1):9-25.

3. Food and Agriculture Organization and World Health Organization Expert Consultation. Evaluation of health and nutritional properties of powder milk and live lactic acid bacteria. Córdoba, Argentina: Food and Agriculture Organization of the United Nations and World Health Organization; 2001.

4. Cave N. Nutritional management of gastrointestinal diseases. In A. J. Fascetti & S. J. Delaney (Eds.), *Applied Veterinary Clinical Nutrition*. Wiley-Blackwell; 2012, 175-219.

5. Satyaraj E, Reynolds A, Pelker R, et al. Supplementation of diets with bovine colostrum influences immune function in dogs. *Br J Nutr*. 2013;110(12):2216-21.

6. Gore AM, Satyaraj E, Labuda J, et al. Supplementation of diets with bovine colostrum influences immune and gut function in kittens. Nestlé Purina PetCare Internal Data.